

# Study of Sorption Isotherms and The Development of Mathematical Model for Varied Temperatures of Cocoyam and Bitter Kola

<sup>1</sup>Iji, Sunday Ibe, <sup>2</sup>Oboh, Innocent Oseriboh, <sup>3</sup>Ekpenyong, Abasiyake Uku

<sup>1</sup>Skills Training Centre, Federal Polytechnic, Ukana. Akwa Ibom State, Nigeria.

<sup>2</sup>Department of Chemical and Petroleum Engineering, University of Uyo, Uyo. Akwa Ibom State, Nigeria.

<sup>3</sup>Department of Chemical and Petroleum Engineering, University of Port Harcourt. Rivers State, Nigeria.

## ABSTRACT

The proximate properties and sorption isotherms of Cocoyam (Taro) and Bitter kola (*Garcinia kola*) have been studied. Cocoyam was processed into pellets of 2 mm while bitter kola was processed into powder of 500  $\mu\text{m}$ . The proximate properties determined were moisture content, crude protein, crude fiber, fat, ash and carbohydrate content. The sorption isotherms were determined using the static gravimetric method. The obtained data were fitted into the Brunauer-Emmett-Teller, Guggenheim-Anderson-de Boer and Oswin models using the nonlinear regression method. From the results obtained, equilibrium moisture content increased with increasing water activity at a particular temperature and decreased with increase in the temperatures studied. The water surface areas of the adsorbents were calculated and the values obtained were less than 260  $\text{m}^2\cdot\text{g}^{-1}$ . The sorption isotherm curves obtained were sigmoidal and the models were also suitable for describing the experimental data with coefficient of determination values of approximately equal to unity. A mathematical model was developed and was fitted to the experimental data which gave the correlation coefficient of  $R^2 = 0.988912$  for taro and  $R^2 = 0.984992$  for *Garcinia kola*. The result of this work revealed that shelf life of cocoyam and bitter kola would be sustained by storage under lower temperatures.

**Keywords:** Bitter kola, Cocoyam, Proximate properties, Sorption Isotherms, Static gravimetric method.

Date of Submission: 05-02-2026

Date of acceptance: 16-02-2026

## I. INTRODUCTION

### 1.1 Background of the Study

Cocoyam is the third most important root crop after yam and cassava in West Africa. It is a predominant food for millions of people living in the tropics. The corms and cormels are rich in minerals, vitamins and digestive starch. According to Iji *et al.* (2021b), about 30-40 species of cocoyam have been identified but only 5 to 6 species produce edible parts. Two genera of cocoyam are widely cultivated in Africa. These are namely *taro* (*Colocasia esculenta*) and *tannia* (*Xanthosoma sagittifolium*).

However, there is a dearth of research and development on cocoyam in Nigeria when compared with other tropical root crops like yam and cassava. Among the reasons for the underutilization of cocoyam is the presence of calcium oxalate raphide; the irritant which causes itching effect felt throughout the throat when consumed. Another reason is that cocoyam is prone to pre-harvest and post-harvest diseases, which reduce storage stability and quality of the tubers. To overcome these challenges, the corms and cormels may be processed into flour. Though cocoyam flour stores better than the corm due to highly reduced moisture content, conversion into flour is not an end to the storage problem (Sunday *et al.*, 2023). According to Obiegbuna *et al.* (2014), all food products are inherently unstable and quality retention depends upon a number of factors including storage temperature, storage relative humidity and storage time.

Bitter kola which is also known as '*Garcinia kola*' is a species of flowering plant in the Clusiaceae or Guttiferae family. It is found in Benin, Cameroon, Democratic Republic of the Congo, Ivory Coast, Gabon, Ghana,

Liberia, Nigeria, Senegal and Sierra Leone (Iji *et al.*, 2021d). Its natural habitat is subtropical or tropical moist lowland forests. The fruit, seeds, nuts, stem and bark of the plant have been used for centuries in folk medicine to treat ailments from coughs to fever, for dental care in the form of chewing-sticks (Iji *et al.*, 2021c). It is highly used for its medicinal purposes because of its anti-viral, anti-inflammatory, anti-diabetic, bronchio-dilator and anti-hepatotoxic attributes. Fruit extracts from it have proven effective at stopping Ebola virus replication in laboratory test (Okunji *et al.*, 2002). It has been utilized in folklore remedies by the Igbo people of the Southern Nigeria to treat ailments associated with poisoning, liver disorders, hepatitis, diarrhoea, laryngitis, bronchitis and gonorrhoea (Iji *et al.*, 2021a).

Some of the use of bitter kola depends on principles of adsorption. Bitter kola is also rich in caffeine and theobromine and is also believed to be an aphrodisiac. Unlike other kola nuts however, bitter kola is believed to clean the digestive system, without side effects such as abdominal problems, even when a lot of nuts are eaten (Iji *et al.*, 2021a).

Adsorption is a mass-transfer surface process in which molecules, ions, or gases accumulate at the interface of a solid adsorbent, driven by attractive forces until dynamic equilibrium is reached when there is no further net transfer (Sunday *et al.*, 2025). The food sorption isotherm describes the thermodynamic relationship between water activity and the equilibrium of moisture content at constant temperature and pressure. The moisture sorption isotherms are extremely important in modeling the drying process, design and optimization of drying equipment, predicting shelf-life stability, calculating moisture changes which may occur during storage and selection of appropriate packaging material. Also, the knowledge of the sorption data is essentially useful to predict microbiological, enzymatic and chemical stability of food materials (Oyerinde and Lawal, 2015). Experimental determination and modeling of sorption isotherms of food materials have attracted numerous researchers because their values are used in industrial purposes. Sorption isotherms are usually classified according to their shape in five different types. The sorption isotherms for most foods are nonlinear and generally with sigmoid shape (Ricardo *et al.*, 2011).

In this current study, sorption properties and the development of mathematical model for varied temperatures of cocoyam and bitter kola are reported. This is expected to provide information for those involved in their processing and application in food and drug formulation.

## II. MATERIALS AND METHODS

### 2.1 Experimental Methods

### 2.2 Preparation of the Raw Materials


Two materials were selected for use in this study: *Taro* and *Garcinia kola*. These were obtained from the Botany Department of University of Uyo, Uyo and are identified as such. They were washed in water, peeled manually with a steel knife, sliced and dried using a hot air oven at 105 °C until constant weight (Iji *et al.*, 2021b and Iji *et al.*, 2021c).

#### 2.2.1 Proximate Analysis

Moisture, crude protein, fibre, fat, lipid, ash and carbohydrate contents were determined according to Association Official of Analytical Chemists (Owuamanam *et al.*, 2010 and Nwanekezi *et al.*, 2010).

### 2.3 Fabricated Air Tight Hermetic Container

The container served as an air tight apparatus that can carry the raw materials and salt solution. The fabrication involved the use of eighteen (18) plastics with cylindrical shape, polyvinyl chloride pipes (2 cm diameter) and aluminum crucibles, cut into circular shape of 4 cm diameter and a lid to cover the container. The design is as shown by Figure 1.0.



**Figure 1.0: Design of the hermetic container**  
Source: The Researcher (2025)

#### 2.4 Preparation of Saturated Salt Solution

Nine saturated saline solutions which include; Lithium Chloride (LiCl), Sodium Chloride (NaCl), Magnesium Chloride ( $MgCl_2$ ), Sodium Bromide (NaBr), Strontium Chloride ( $SrCl_2$ ), Potassium Chloride (KCl), Potassium Carbonate ( $K_2CO_3$ ), Potassium Iodide (KI) and Potassium Sulphate ( $K_2SO_4$ ) were used, these were enclosed in hermetic containers and had  $a_w$  values that varied between 0.113 and 0.970 (Iji *et al.*, 2021a; Iji *et al.*, 2021b; Iji *et al.*, 2021c; Iji *et al.*, 2021). 3g of *taro* and *garcina kola* were introduced into each hermetic container with the corresponding saline solution. Each analytical grade salt was mixed with distilled water (approximately 100 ml) until a solution with excess crystals was formed. The saturated salt solutions were poured into the hermetic containers to a depth of 1 cm (Iji *et al.*, 2021a).

The adsorption isotherms were determined using a temperature-controlled cabinet (Hot Pack, US); with an accuracy of  $\pm 1$  °C at the selected temperatures of 30 °C to 60 °C using the static gravimetric method, according to the COST 90 project. Thymol was placed into the containers with saline solutions that had  $a_w$  values higher than 0.65 to prevent microbial growth. The samples were periodically weighed until a constant weight was achieved, to ensure equilibrium between the samples and the saline solutions. The adsorption experiments for each temperature were performed in triplicate (Iji *et al.*, 2021a).

#### 2.5 Sorption Isotherm

The relationship between the equilibrium moisture content and the water activity of the product was predicted by using the equations representing the models commonly used in foodstuff. These models which incorporate the temperature effect have been adopted as standard equations by the ASAE for the description of sorption isotherms (Iji *et al.*, 2021a). In this study, the three isotherm models used to fit the experimental data are presented in Table 1.

Table 1: Isotherm models and their equations

Name of Model	Range of $a_w$	Mathematical Expression
BET (Brunauer-Emmett-Teller)	$0.05 < a_w < 0.35$	$\frac{M}{M_0} = \frac{C a_w}{(1 - a_w)(1 - a_w + C a_w)}$
GAB (Guggenheim-Anderson-de Boer)	$a_w < 0.95$	$M = \frac{M_0 G K a_w}{(1 - K a_w)(1 - K a_w + G K a_w)}$
Oswin	$0.10 < a_w < 0.80$	$M = A \left[ \frac{a_w}{1 - a_w} \right]^B$

Source: Tabulated by researcher (2021)

The constants were estimated by fitting the mathematical model to the experimental data, using a non-linear regression analysis with Microsoft Excel 2013 software.

The quality of the fitting of different models was evaluated by calculating the correlation coefficient ( $R^2$ ), the mean relative percentage deviation modulus E in percentage (%) and the root mean square error (RMSE) between the experimental and predicted equilibrium moisture content (Basu *et al.*, 2006). The deviation modulus and root mean square error and correlation coefficient are expressed as Equations 1 through 3.

$$E = \frac{100}{N} \sum_{i=1}^N \frac{|X_{eq,exp} - X_{eq,pre}|}{X_{eq,exp}} \tag{Equation 1}$$

$$RMSE = \left[ \frac{1}{N} \sum_{i=1}^N \frac{(X_{eq,exp} - X_{eq,pre})^2}{X_{eq,exp}} \right]^{\frac{1}{2}} \tag{Equation 2}$$

$$R^2 = \frac{(X_{eq,exp})}{X_{eq,exp} + X_{eq,pre}} \tag{Equation 3}$$

Where N is the number of observations;  $X_{eq,exp}$  and  $X_{eq,pre}$  are the experimental and predicted values of the equilibrium moisture content, respectively.

## 2.6 Surface area

The specific surface area plays an important role in determining the water binding properties of particulate products (Iji *et al.*, 2021a). It is represented by Equation 4.

$$S_a = X_m \frac{1}{M} N_A = 3.53 \times 10^3 X_m \tag{Equation 4}$$

Where  $S_a$  is the product surface area ( $m^2 \cdot g^{-1}$ ),  $X_m$  is the monolayer moisture content ( $kg \cdot kg^{-1}$  dry basis), M is the molecular weight of water ( $18 g \cdot mol^{-1}$ ),  $N_A$  is the Avogadro's number ( $6 \times 10^{23}$  molecules  $\cdot mol^{-1}$ ) and A is the area of one water molecule ( $1.06 \times 10^{-19} m^2$ ). Where  $S_a$  is the product surface area ( $m^2 \cdot g^{-1}$ ),  $X_m$  is the monolayer moisture content ( $kg \cdot kg^{-1}$  dry basis), M is the molecular weight of water ( $18 g \cdot mol^{-1}$ ),  $N_A$  is the Avogadro's number ( $6 \times 10^{23}$  molecules  $\cdot mol^{-1}$ ) and A is the area of one water molecule ( $1.06 \times 10^{-19} m^2$ ).

## 2.7 Development of Mathematical Models

Literatures had established that the non – linear regression (or equation) technique could be used to develop a model which takes care of many variables (Khazei and Daneshmandi, 2007; Maisnam *et al.*, 2017; Iji *et al.*, 2021a).

For the development of the correlation in this study, The BET model equation was considered as shown in Table 1. The Moisture content (M) depends directly on the monolayer moisture content ( $M_0$ ), net heat of sorption (C) and water activity ( $a_w$ ). Based on this assertion, the relationships of M with the adsorption parameters are presented in Equations 5 through 7.

$$M \propto M_0 \tag{Equation 5}$$

$$M \propto C \tag{Equation 6}$$

$$M \propto a_w \tag{Equation 7}$$

For the development of the proposed moisture ratio correlation for adsorption of the products (bitter kola and cocoyam), the following parameters, namely: monolayer moisture content, net heat of sorption and water activity. In other words, the moisture ratio is expressed as a function of the mentioned parameters in Equations 8.

$$M = f(M_0, C, a_w) \tag{Equation 8}$$

Equations 5 through 7 can be expanded in the basic form of nonlinear multivariable algebraic expression as given in Equation 9.

$$M = \beta_0 (M_0^{\beta_1} \cdot C^{\beta_2} \cdot a_w^{\beta_3}) \tag{Equation 9}$$

Alternatively, Equation 14 can also be expressed as Equation 10.

$$M = \beta_o [M_o^{\beta_1} \times C^{\beta_2} \times a_w^{\beta_3}] \quad \text{Equation 10}$$

Where  $\beta_o$  is the adsorption coefficient while  $\beta_1$ ,  $\beta_2$  and  $\beta_3$  are exponential constants of the drying parameters.

The variables  $\beta_o$ ,  $\beta_1$ ,  $\beta_2$  and  $\beta_3$  could be determined iteratively using multivariable numerical optimization method. To determine these variables, existing literature indicates that, there are several approaches to performing this iterative estimation, namely; Gauss-Newton, the Marquardt-Levenberg, the Nelder-Mead, the steepest descent, General Reduced Gradient (GRG) protocol (Wei, 2013; Smyth, 2015; Pashaie *et al.*, 2016; Okon *et al.*, 2015).

To estimate the variables in the Equation 15 iteratively, these parameters were obtained experimentally. Then the unknown coefficient ( $\beta_o$ ) and the exponential constants  $\beta_1$ ,  $\beta_2$  and  $\beta_3$  were solved iteratively using GRG protocol in the Microsoft Excel SOLVER. To validate the correlation, its prediction was compared with the experimental data and some authors' correlation predictions. Also, the closeness of the correlation's predictions and experimental data were established using coefficient of determination ( $R^2$ ) and other error functions.

### III. RESULTS AND DISCUSSION

#### 3.1 Results and Discussion

##### 3.1.1 Proximate Analysis of *Taro* and *Garcina Kola*

Table 2 shows the moisture, crude protein, crude fibre, fat, ash and carbohydrate contents of *taro* and *garcina kola*. These materials possess high moisture content, a property of any food or fruit that can make it viable to microbial attack and this accounts for most biochemical and physiological reactions in plant foods. The high protein content in *taro* is desirable on nutritional grounds (Nwankezi *et al.*, 2010). From the value obtained, it can be deduced that *taro* is highly nutritional.

Fibre plays a more important role in human nutrition than its functionality in the food system. It contributes to bowel movement and cocoyam has been a health food for people with gastro intestinal disorders (Iji *et al.*, 2021a). This implies that *garcina kola* will serve better for this purpose.

The carbohydrate content for *taro* is high and doesn't differ among the cultivars thus signifying the importance of cocoyam as energy giving food nutrient.

**Table 2: Proximate composition for *taro* and *garcina kola***

Components	Percentage (%)	
	<i>Taro</i>	<i>Garcina Kola</i>
Moisture content	70.300	52.500
Crude protein	3.210	2.510
Crude fibre	0.860	5.120
Fat content	1.400	4.160
Ash	2.150	0.860
Carbohydrate	22.080	34.850

Source: Computed by the researcher (2021)

##### 3.2 Equilibrium moisture content and water activities at varied temperatures

The results of the EMC and  $a_w$  of *taro* and *garcina kola* are shown in Tables 3 and 4. These were taken at temperature range of 30 to 60 °C. It can be clearly observed that as temperature increases, the EMC decreases for both materials.

**Table 3: Summary of the Equilibrium Moisture Content and Water Activity at Varied Temperatures for *Taro***

<i>Taro</i>							
30 °C		40 °C		50 °C		60 °C	
$a_w$	EMC	$a_w$	EMC	$a_w$	EMC	$a_w$	EMC
0.113	0.010	0.112	0.007	0.111	0.005	0.111	0.004
0.324	0.015	0.316	0.010	0.305	0.007	0.293	0.006
0.432	0.018	0.432	0.013	0.432	0.009	0.432	0.008
0.560	0.027	0.532	0.024	0.509	0.020	0.497	0.018
0.679	0.052	0.661	0.048	0.645	0.044	0.631	0.041
0.691	0.053	0.673	0.049	0.657	0.045	0.643	0.043

0.751	0.062	0.747	0.058	0.744	0.052	0.745	0.049
0.836	0.117	0.823	0.113	0.812	0.111	0.803	0.109
0.970	0.700	0.964	0.570	0.958	0.483	0.952	0.409

Source: Computed by the researcher (2021)

Table 4: Summary of the Equilibrium Moisture Content and Water Activity at Varied Temperatures for *Garcina Kola*

30 °C		40 °C		50 °C		60 °C	
a <sub>w</sub>	EMC						
0.113	0.050	0.112	0.049	0.111	0.048	0.111	0.045
0.324	0.070	0.316	0.068	0.305	0.066	0.293	0.065
0.432	0.081	0.432	0.080	0.432	0.079	0.432	0.077
0.560	0.110	0.532	0.100	0.509	0.097	0.497	0.095
0.679	0.112	0.661	0.120	0.645	0.117	0.631	0.114
0.691	0.127	0.673	0.125	0.657	0.120	0.643	0.118
0.751	0.200	0.747	0.199	0.744	0.197	0.745	0.190
0.836	0.330	0.823	0.310	0.812	0.290	0.803	0.285
0.970	1.835	0.964	1.830	0.958	1.790	0.952	1.750

Source: Computed by the researcher (2021)

### 3.3 Moisture Sorption Isotherm of *Taro* and *Garcina Kola* at Varied Temperatures

The experimental data for the EMC and a<sub>w</sub> were plugged into nonlinear regression graphs using Microsoft Excel 2013. Figures 2 through 5 are the moisture sorption isotherms for *taro* and *garcina kola* at varied temperatures. The equilibrium moisture at a given water activity, decreased with increase in temperature. This is in agreement with the theory of physical sorption (Iglesias *et al.*, 1975; Hassian *et al.*, 2001). Implying that *taro* and *garcina kola* became less hygroscopic with increasing temperature. The adsorption isotherms exhibited Type II curve with a characteristic sigmoidal shape.




Figure 2: Adsorption isotherm of *taro* and *garcina kola* at 30 °C  
Source: Plotted by the researcher (2021).




Figure 3: Adsorption isotherm of *taro* and *garcina kola* at 40 °C  
Source: Plotted by the researcher (2021).




Figure 4: Adsorption isotherm of *taro* and *garcina kola* at 50 °C  
Source: Plotted by the researcher (2021).




Figure 5: Adsorption isotherm of *taro* and *garcina kola* at 60 °C  
Source: Plotted by the researcher (2021).

### 3.4 Adsorption Models

The experimental data (EMC) obtained were plugged into the adsorption models which include Brunauer-Emmett-Teller (BET), Guggenheim-Anderson-de Boer (GAB) and Oswin models. Tables 5 and 6 show the coefficients of the models, the mean relative percentage modulus (E), the root mean square error (R.M.S.E) and the correlation coefficient ( $R^2$ ). All the models presented correlation coefficients very close to unity indicating good fit to experimental data. The high values of  $R^2$  at different temperatures revealed that the good fit of the models is independent of temperature variation.

**Table 5: Estimated parameters by Non-linear Regression for BET, GAB and OSWIN Models for *Taro***

Models	Constants	Temperatures (°C)			
		30	40	50	60
BET	$M_0$ (g/g dry solid)	0.021510	0.021155	0.021111	0.020672
	C	1.291558	1.284853	1.182593	1.206428
	$R^2$	0.999672003	0.999034687	0.997019764	0.993727462
	RMSE	0.003786	0.005265	0.007858	0.009655
	E (%)	7.517906	1.07209	13.1379	19.396
GAB	$M_0$ (g/g dry solid)	0.02872	0.027912	0.02467	0.024434
	K	0.9913	0.99025	0.877883	0.989999
	G	1.210937	1.090866	1.154837	1.150999
	$R^2$	0.994268	0.995156	0.997222	0.993003
	RMSE	0.0016255	0.0012121	0.007544	0.00989
E (%)	14.9403	19.5608	16.0394	33.1579	
OSWIN	A	0.024294	0.024151	0.023377	0.023377
	B	0.971778	0.962395	0.962395	0.961963
	$R^2$	0.999258	0.999171	0.996423	0.998151
	RMSE	0.005787	0.004874	0.004874	0.008363
	E (%)	6.461344	0.47165	0.47165	15.1069

Source: Computed by the researcher from calculated data (2021)

**Table 6: Estimated Parameters by Nonlinear Regression for BET, GAB and OSWIN Models for *Garcinia Kola***

Models	Constants	Temperatures (°C)			
		30	40	50	60
BET	$M_0$ (g/g dry solid)	0.181576	0.080867	0.041238	0.032293
	C	0.552673	0.917647	1.386102	1.706528
	$R^2$	0.99835	0.998261	0.997453	0.993035
	RMSE	0.02211	0.022701	0.026876	0.045547
	E (%)	12.56221	16.31022	17.69961	10.38048
GAB	$M_0$ (g/g dry solid)	0.072897	0.050659	0.029747	0.023466
	K	0.989682	0.999365	0.974841	1.007904
	G	1.002062	1.174062	1.666545	1.796874
	$R^2$	0.996416	0.998076	0.997479	0.997517
	RMSE	0.032727	0.023936	0.026832	0.026348
E (%)	16.50219	21.07163	20.83303	25.12812	
OSWIN	A	0.079592	0.068519	0.065333	0.062802
	B	0.903281	0.999387	1.058778	1.114019
	$R^2$	0.99678	0.998234	0.99791	0.997649
	RMSE	0.030881	0.022931	0.02446	0.025406
	E (%)	11.87691	18.26048	21.06275	23.0462

Source: Computed by the researcher from calculated data (2021)

### 3.5 Surface Area

Table 7 presents the surface area of *taro* and *garcina kola* from 30 to 60°C. It was estimated using Equation 4 and the monolayer moisture contents of *taro* and *garcina kola* were obtained by GAB model. The results indicated that, the availability of surface areas for hydrophilic binding decrease with increasing temperature. The calculated surface area values of *taro* and *garcina kola* were within the range commonly obtained for food stuffs (100 - 250 m<sup>2</sup>.g<sup>-1</sup>) (Cassini *et al.*, 2006).

**Table 7: Surface area of *taro* and *garcina kola***

Temperature (°C)	Adsorption S <sub>a</sub> (m <sup>2</sup> .g <sup>-1</sup> )	
	<i>Taro</i>	<i>Garcina Kola</i>
30	101.38180	257.32650
40	98.52780	178.82640
50	87.08450	105.00690
60	86.25210	82.83500

Source: Computed by the researcher (2021)

### 3.5 Developed Models

Three parameters were used to develop a mathematical model that relates moisture content (M) to monolayer moisture content (M<sub>0</sub>), net heat of sorption (C) and water activity (a<sub>w</sub>) for the adsorption of moisture by *taro* pellet and *garcina kola* powder to which this research is established. Table 8 shows the values of the developed adsorption coefficient (β<sub>0</sub>) and exponential constants (β<sub>1</sub>, β<sub>2</sub>, and β<sub>3</sub>). These values were obtained using Microsoft Excel 2013 Solver after iteration with the General Reduced Gradient (GRG) protocol.

**Table 8: Simulated values for different product using the developed model**

Symbol	<i>Taro</i>	<i>Garcina Kola</i>
β <sub>0</sub>	104.70550	96.69954
β <sub>1</sub>	1.51878	1.80472
β <sub>2</sub>	0.19998	0.51181
β <sub>3</sub>	10.58964	10.09131

Source: Computed by the researcher (2021)

The quality of the developed model was evaluated using different statistical criteria to check for goodness of fit. The statistical variables include; correlation coefficient (R<sup>2</sup>), root mean square error (RMSE) and sum square error (SSE). This is as shown in Table 9.

**Table 9: Statistical values for different product using the developed model**

Symbol	<i>Taro</i>	<i>Garcina Kola</i>
R <sup>2</sup>	0.98891	0.98499
RMSE	0.02297	0.07039
SSE	4.83446	4.63077

Source: Computed by the researcher (2021)

The empirical model developed is given as Equation 11 and Equation 12 for *taro* and *garcina kola* respectively.

$$M = 104.70550[M_0^{1.51878} \times C^{0.19998} \times a_w^{10.58964}] \quad \text{Equation 11}$$

$$M = 96.69954[M_0^{1.80472} \times C^{0.51181} \times a_w^{10.09131}] \quad \text{Equation 12}$$

## IV. Conclusion

This study has specifically provided the following information on sorption properties for varied temperature of cocoyam (*taro*) pellets bitter kola (*garcina kola*), and the following had been revealed:

- i. The proximate properties and sorption isotherms of *garcina kola* powder and *taro* pellets were determined.

- ii. The equilibrium moisture content of *taro* and *garcina kola* decreased with increase in temperature at a given water activity ( $a_w$ ), indicating that they become less hygroscopic at high temperatures giving a clear stability domain of *taro* and *garcina kola* after drying leading to extensive shelf life.
- iii. The results of the fit revealed that the three models accurately described all the adsorption and moisture isotherms for *taro* and *garcina kola*.
- iv. The mathematical models were developed and validated by fitting it into the experimental data which produced accurate results of correlation coefficient ( $R^2 = 0.988912$  for *taro* and  $R^2 = 0.984992$  for *garcina kola*).

### References

- [1]. Basu, S., Shivhare, U. S. and Mujumdar, A. S. (2006). Models for sorption isotherms for foods: A review. *Dry Technology*, 24(8): 917 – 930.
- [2]. Cassini, A. S., Marczak, L. D. F. and Norena, C. P. Z. (2006). Water adsorption isotherms of texturized soy protein. *Journal of Food Engineering*, 77: 194–199.
- [3]. Catherine, B., Paul, L., Claire, R., and Caetano, C. D. (2019). Experimental determination of moisture sorption isotherm of fecal sludge, vol. 11, pp. 1 - 7; doi:10.3390/w11020303.
- [4]. Hahn, S. K., Osiru, D. S. O., Akoroda, M. O. and Otoo, J. A. (1987). Yam production and its future prospects In: Outlook on Agriculture, IITA, Ibadan, 16: 105-108.
- [5]. Hassian, M. D., Bala, B. K., Hossian, M. A. and Mondolo, M. R. A. (2001). Sorption isotherm and isosteric heat of sorption of pineapple. *Journal of Food Engineering*, 48, 103-107.
- [6]. Iglesias, H. A., Chirife, J., and Lombardi, J. L. (1975). An equation for correlating equilibrium moisture content in foods, *Journal of Food Technology*, 10: 289–294.
- [7]. Jji, S. I., Oboh, I. O. and Ekpenyong, A. U. (2021a). Sorption properties and the development of mathematical model for varied temperatures of bitter kola, *International Journal of Engineering and Technology Research*, 22(5): 1-14.
- [8]. Jji, S. I., Oboh, I. O. and Ekpenyong, A. U. (2021b). The study of sorption isotherms for varied temperatures of cocoyam (*taro*), *Journal of Management Science and Entrepreneurship*, 19(7): 301-316.
- [9]. Jji, S. I., Oboh, I. O. and Obomanu, C. A. (2021c). The study of sorption isotherms for varied temperatures of bitter kola (*garcina kola*), *International Journal of Pure and Applied Science*, 20(9): 268-281.
- [10]. Jji, S. I., Oboh, I. O. and Obomanu, C. A. (2021d). The study of sorption isotherms for varied temperatures of bitter kola (*garcina kola*), *Journal of African Sustainable Development*, 19(2): 242-256.
- [11]. Khazaei, J. and Daneshmandi, S. (2007): Modelling of thin-layer drying of sesame seeds: Mathematical and neural networks modeling, *International Agrophysics*, 21:335 - 348.
- [12]. Nwanekezi, E. C., Owuamanam, C. I., Ihediohanma, N. C., and Iwouno, J. O. (2010). Functional, particle size and sorption isotherm of cocoyam cormel flours. *Pakistan Journal of Nutrition*, 9 (10): 973-979.
- [13]. Obiegbona, J. E., Ishiwu, C. N., Akubor, P. I. and Igwe, E. C. (2014). Effect of processing and storage relative humidity on selected functional properties of cocoyam (*colocasia esculenta*) corm flour, *Journal of Food Science and Quality Management*, 28:19-28.
- [14]. Okon, A. N., Udoh, F. D. and Appah, D. (2015). Empirical Wellhead Pressure - Production Rate Correlations for Niger Delta Oil Wells. *Paper presented at the Society of Petroleum Engineers Nigeria Annual International Conference and Exhibition*, Lagos, Nigeria, 4-6 August, pp. 1-17.
- [15]. Okunji, C. O., Ware, T. A., Hicks R. P., Iwu, M. M., Skanchy, D. J. (2002). Capillary electrophoresis determination of biflavonones from *Garcinia kola* in three traditional African medicinal formulations. *Planta Med.*, 68, pp. 440 – 444.
- [16]. Onwueme, I. C. and Simha, T. D. (1991). Field crop production in tropical Africa. CTA Ed, Netherlands, pp: 276-288.
- [17]. Owuamanam, C. I., Ihediohanma, N. C. and Nwanekezi E. C. (2010). Sorption isotherm, particle size, chemical and physical properties of cocoyam corm flours. Department of Food Science and Technology Federal University of Technology, PMB 1526, Owerri, Nigeria, pp. 11-17.
- [18]. Pashaie, M., Sadeghi, M. and Jafarian, A. (2016). Artificial Neural Networks with Nelder-Mead Optimization Method for solving Nonlinear Integral Equations. *Journal of Computer Science and Application*, 8(1): 1-20
- [19]. Ricardo, D., Andrade P., Roberto, M., Carmen E. and Pérez C. (2011). Models of sorption isotherms for food: Uses and Limitation, pp. 325-334.
- [20]. Smyth, G. K. (2015). Optimization and Nonlinear Equations. Wiley StataRef: Statistics Reference Online. pp. 1-9, doi: 10.1002/9781118445112.stat05030.pub2.
- [21]. Sunday, I. I., Abasiyake, U. E. and Richard, U. (2023). Academic Staff Union of Polytechnic (ASUP) (Zone D), 6<sup>th</sup> Natural Conference, Kenule Benson Saro-Wiwa Polytechnic, Bori, Port Harcourt, pp. 1-15.
- [22]. Sunday, I. I., Uwem, E. I. and Benjamin, R. E. (2025). Models for the development of sorption isotherms: A review. *American Journal of Food Science and Technology*, 13(2): 27-37.